



+WEEDKILL

SLOW RELEASE LAWN FERTILISER

This product is specially formulated to provide deep uniform greening, whilst killing most common broadleaf weeds. The slow release formula feeds the lawn as it needs to be fed, avoiding surge growth and extra mowing.



+WEEDKILL

SLOW RELEASE LAWN FERTILISER

GUARANTEED ANALYSIS

Scotts Lawn Builder NPK 20:1:15 (N : P=20)

20%	Nitrogen (N)	20% as slow release methylene urea and free urea
1.0%	Phosphorus (P)	water soluble
15%	Potassium (K)	as potassium sulphate
7.5%	Sulphur (S)	as sulphate
2%	Iron (Fe)	(20000mg/kg)

This product contains below detectable levels of cadmium (Cd), lead (Pb) and mercury (Hg). Therefore this product may be used without restriction on food and animal feed crops.

STORAGE AND DISPOSAL

Store in the closed original container in a cool, dry place out of the reach of children. Do NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in the garbage. Sweep up spilled material and return to original container. Spoilt product should be disposed of as should the empty container.

SAFETY DIRECTIONS

Will irritate the eyes. Avoid contact with eyes. When opening the container and using the product, wear elbow-length PVC gloves and goggles and disposable dust mask covering mouth and nose. Wash hands after use. After each day's use, wash gloves and goggles.

FIRST AID

If poisoning occurs, contact a doctor or poisons information centre. Phone Australia 131 126 or New Zealand 0800 764 766. For specialist advice in an emergency only, phone NZ 0800 734 607. If skin contact occurs, remove contaminated clothing and wash skin thoroughly. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

FOR FREE GARDENING TIPS AND INFORMATION,
CALL OUR GARDEN HELPLINE TOLL FREE:

Australia 1800 804 219 or
NZ 0800 449 213

www.scottsaustralia.com.au



DIRECTIONS FOR USE

IMPORTANT Do not use on Buffalo grass, Carpet grass, Santa Ana Couch, Lippia or Dichondra as damage may occur.

LAWN TYPE	WEED CONTROLLED
Suitable for lawns containing common Couch, Bent (not Bent greens), Kikuyu, Perennial Ryegrass and/ or Chewings Fescue only.	Broadleaf weeds including Bindii (Onehunga Weed), Catsear (Flatweed), Chickweed, Mouse-eared Chickweed, Clover, Dandelion, Fat Hen, Creeping Oxalis, Mallow, Pearlwort, Plantain (Lambs Tongue), Staggerweed, Fumitory, Cobblers Peg, Hydrocotyl (Pennyweed), Capeweed and Wireweed.

HOW DO I APPLY IT?

- This product is ready to apply. Apply evenly at the rate of 16g/m².
- Apply to moist foliage. If foliage is not moist from dew or recent rain, wet the area to be treated. Leave on foliage for 24 hours before watering in thoroughly.
- For best results use a Scotts Spreader, however, if applying by hand, always wear gloves and spread evenly moving in a steady forward motion.

Do not attempt to spread the product in one pass – you may under or over apply. Apply by passing over the treated area at least three times, moving in a different direction each time and applying about a third of the required amount each pass.

- Rain, watering or heavy traffic within 24 hours can knock granules off weeds and may reduce effectiveness. Do not mow within 48 hours of applying this product.
- Re-treating may be necessary, but wait at least three weeks to observe the full results of the first application. To obtain even greening and weed control it is essential that the granules be applied evenly. Poor or no weed control will occur in untreated patches. Apply under calm conditions. Windy conditions will make even distribution difficult.

APPLYING BY SCOTTS SPREADERS



EvenGreen Drop Spreader
- SETTING 5



EasyGreen Rotary Spreader
- SETTING 26



HandyGreen Hand Held Spreader
- SETTING 4¼

IMPORTANT: THIS PRODUCT IS NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED. See CAUTION statements for essential information about use of this product.

When to Apply

Best control will be obtained when weeds are actively growing and young, rather than flowering or seeding. Do not apply if weeds are stressed due to dry, frosty or excessively moist conditions. Do not apply if rain or irrigation is expected within 24 hours. Avoid application when daytime temperatures are consistently above 30°C.

IMPORTANT

This product when wet, will cause iron stains. Be sure to sweep all granules off paths, coping, paving, driveways, shoes, etc. immediately after use. Do not allow the granules to come into contact with these surfaces when wet.

PROFESSIONAL QUALITY GUARANTEE



The Scotts Difference

Unlike ordinary weed 'n' feeds, every particle of Scotts Lawn Builder™ with Weed Kill contains an effective triple action weed control agent and balanced nutrient mix, including Phosphorus, Potassium and Scotts patented slow release Nitrogen. With correct application, this product is guaranteed to provide consistent, professional quality results at home.



Scotts Australia Pty. Ltd.
ACN 003 123 162
Building E, Level 2, 32 Lexington Drive,
Bella Vista NSW 2153 Australia, or
L4, 152 Fanshawe St. Auckland, New Zealand.

APVMA No. 49752/ 0607
HSNO Approval Code HSR000365 (Dicot III).
ACVM approval not required.




+WEEDKILL

SLOW RELEASE LAWN FERTILISER



Clover




Bindii



Creeping Oxalis



Dandelion



Lambs Tongue

RECOMMENDED APPLICATION TIMES

WEED	WHEN TO APPLY
Bindii (Onehunga Weed), Capeweed, Catsear (Flatweed), Chickweed, Dandelion, Fumitory, Hydrocotyl (Pennyweed), Mallow, Mouse-eared Chickweed, Plantain (Lambs Tongue)	Autumn to early Spring
Cobblers Peg, Creeping Oxalis, Chickweed, Dandelion, Hydrocotyl (Pennyweed), Plantain, Fat Hen, Catsear (Flatweed)	Spring & Summer
Clover, Pearlwort, Staggerweed, Wireweed	Year round

CAUTION

- On new grass plantings wait 4 weeks after the first mowing before application.
- Do not plant new grass within one month of application.
- Do not mow grass within 48 hours of application.
- Do not apply when daytime temperatures are expected to exceed 30°C.
- Do not over apply, especially under the branch spread of trees and shrubs.
- Do not apply on or around food /animal feed crops

WARNING

- Do NOT allow granules to come in contact with or drift onto your plants you do not want killed, such as shrubs, flower beds and vegetable gardens. Grapevines and tomatoes are particularly sensitive to this product.
- Do NOT allow containers or granules to get into drains, sewers, streams or ponds.
- Do NOT feed grass clippings from treated areas to poultry or other animals.
- Harmful to Aquatic organisms, very toxic to some plant species