

**CAUTION**  
KEEP OUT OF REACH OF CHILDREN

### BORON DEFICIENCY

Boron is important to growing tissue in young shoots, roots, flower buds and fruits. Boron deficiency is more likely on alkaline soils, or those that have been limed heavily.

Deficiency symptoms vary between plant species. Symptoms that may occur include: death of terminal & lateral buds, young leaves thicken, curl & become brittle, lower petals of flowers become brown & eventually die. Young leaves may also show a yellow chlorosis sometimes with reddish-purple colouring. Repeat application every week for 2 to 3 weeks.

### FIRST AID

¥ For advice, contact a Poisons Information Centre (Phone 131 126) or a doctor.

BARCODE  
9315138003122

MEASURING  
SPOON  
INCLUDED

Searles®

**BORAX**

(BORON)

- ✓ Soluble & quick acting
- ✓ Develops growing tissue in young shoots, roots, flower buds and fruits

**NET 500g**

### DIRECTIONS FOR USE

1 HEAPED  TEASPOON = 5g

### EASY GUIDE CHART

APPLICATION	(g)	TEA SPOONS	IN LITRE WATER	COVERAGE
Ground Application	5	1	5Litres	25sq m
Foliar Application	5	1	5Litres	

Spray evenly over foliage.  
DO NOT apply in direct sunlight.

### BORAX ANALYSIS

11% Boron (B) present as Sodium Borate

PACKED BY J.C. & A.T. SEARLE PTY LTD  
4914 D'Aguilar Hwy,  
PO Box 183, Kilcoy QLD 4515 AUSTRALIA  
WEBSITE: [www.searle.com.au](http://www.searle.com.au)  
EMAIL: [searleh@searle.com.au](mailto:searleh@searle.com.au)