

**HOW TO USE: Shake well before use.** • DO NOT spray when shade temperatures exceed 35 °C. • DO NOT spray when soil is dry and plants are suffering from moisture stress. • Applying as a protective spray will give more effective control. • Ensure thorough leaf coverage.

CROP	DISEASE	RATE	HOW TO APPLY
Young or small citrus and recently transplanted trees	Phytophthora Root Rot, Collar Rot	5-10 mL/L	Use higher rate under higher disease risk conditions (e.g. for trees in heavy clay soil). Apply each year as a preventative foliar spray, in late winter prior to flowering and again in Autumn when fruit are approaching maturity. Spray until leaves are just wet. Use 5-10mL/L rate for container grown Mandarin trees. Removal of fruit from affected trees will assist recovery.
Mature citrus (except container grown Mandarin trees)		100-160 mL in 12L water per tree.	
Ornamentals	Phytophthora Root Rot, Collar Rot, Crown Rot	5 mL/L	Apply at 4-6 weekly intervals as foliar spray when conditions favour disease. Certain plants may be sensitive to phosphorous acid sprays. Test on a small sample first if in doubt. DO NOT apply when plants are dormant or stressed.
Avocado	Phytophthora Root Rot	2.5-3 mL/L	Apply as a foliar spray to run-off. Use 7.5-10L per adult tree. Apply every 3 weeks as a curative spray, or every 5-6 weeks as a preventative spray.
Grapes	Downey Mildew	6 mL/L	Apply as a foliar spray to the point of run-off. Apply at first sign of disease.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED.**

**HOW TO PREPARE:**

Mix the required rate with water.

**CAUTION:**

DO NOT allow chemical containers or spray to get into drains, sewers, streams or ponds.

**STORAGE AND DISPOSAL:**

Store in the closed, original container in a cool, dry place out of the reach of children. DO NOT store in direct sunlight. Dispose of empty container by wrapping in paper, placing in a plastic bag and putting in garbage.

**SAFETY DIRECTIONS:**

May irritate the eyes and skin. Avoid contact with eyes and skin. Wash hands after use.

**FIRST AID:**

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 13 11 26.

**MSDS:**

Additional information is listed in the Material Safety Data Sheet which can be obtained from the supplier.

APVMA Approval No: 66554/53859

**DISCLAIMER:**

This product is designed to be used only for the purposes indicated on the label. Failure to use the product strictly as directed may be dangerous and cause injury, may be illegal, and may also render the product ineffective. The buyer or user is responsible for any misuses or negligence in the handling, storage, use or disposal of this material. JC & AT Searle Pty Ltd will not accept any responsibility of any kind, whether for consequential loss or otherwise, in connection with the supply and use of the product other than those responsibilities mandatorily imposed by statutes applicable to the safety and supply of these goods.

Searles®  
**ROOT ROT**  
SYSTEMIC FUNGICIDE

READ SAFETY DIRECTIONS

Searles®  
**ROOT ROT**  
SYSTEMIC FUNGICIDE

ACTIVE CONSTITUENT:  
**200g/L PHOSPHOROUS ACID**  
present as MONO-DI POTASSIUM PHOSPHITE



**Controls Root Rot, Collar Rot, Downey Mildew. For Citrus, Avocadoes, Grapes & Ornamentals.**

The bottle must not be sold separately. Do not destroy box while product still remains.

Contents 500mL



Searles®

**Controls Root Rot, Collar Rot, Downey Mildew. For Citrus, Avocadoes, Grapes & Ornamentals.**



JC & AT Searle Pty Ltd  
4914 D'Aguilar Hwy, KILCOY QLD 4515  
ABN 14 002 898 893  
Phone: (07) 5422 3000  
Fax: (07) 5497 1997

[www.searles.com.au](http://www.searles.com.au)